



## HIV testing and access to HIV medical care among people who inject drugs and their intimate partners in Kazakhstan

Stacey A. Shaw, Assel Terlikbayeva, Leila Famouri, Tim Hunt, Louisa Gilbert, Yelena Rozental, Shopan Primbetova, Mingway Chang, Xin Ma & Nabila El-Bassel

To cite this article: Stacey A. Shaw, Assel Terlikbayeva, Leila Famouri, Tim Hunt, Louisa Gilbert, Yelena Rozental, Shopan Primbetova, Mingway Chang, Xin Ma & Nabila El-Bassel (2016): HIV testing and access to HIV medical care among people who inject drugs and their intimate partners in Kazakhstan, *Journal of Substance Use*, DOI: [10.3109/14659891.2016.1143046](https://doi.org/10.3109/14659891.2016.1143046)

To link to this article: <http://dx.doi.org/10.3109/14659891.2016.1143046>



Published online: 25 May 2016.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

ORIGINAL ARTICLE

## HIV testing and access to HIV medical care among people who inject drugs and their intimate partners in Kazakhstan

Stacey A. Shaw<sup>1</sup>, Assel Terlikbayeva<sup>2</sup>, Leila Famouri<sup>3</sup>, Tim Hunt<sup>1</sup>, Louisa Gilbert<sup>1</sup>, Yelena Rozental<sup>2</sup>, Shopan Primbetova<sup>2</sup>, Mingway Chang<sup>1</sup>, Xin Ma<sup>1</sup>, and Nabila El-Bassel<sup>1</sup>

<sup>1</sup>Global Health Research Center of Central Asia, Columbia University, New York City, New York, NY, USA; <sup>2</sup>Global Health Research Center of Central Asia, Columbia University, Almaty, Kazakhstan; and <sup>3</sup>School of Medicine, West Virginia University, Morgantown, WV, USA

### Abstract

**Background:** Growing rates of HIV and high rates of injection drug use in Kazakhstan call for examining access to testing and treatment among people who inject drugs and their intimate partners.

**Objectives:** We examine how access to health and drug treatment services as well as risk environment factors are associated with ever being tested for HIV and ever receiving any general HIV medical care among 728 male and female intimate partners, where at least one partner injects drugs.

**Methods:** Multivariate random effects logistic regression with random effects for couple were conducted to examine associations between access to health and drug treatment services, risk environment factors, and HIV testing and HIV medical care outcomes.

**Results:** Analyses indicate that accessing needle exchange services and having a regular physician were associated with access to both HIV testing and HIV medical care. Receiving drug treatment was associated with accessing HIV testing but not HIV medical care. Being arrested and charged with a criminal offense was also associated with accessing HIV testing but not HIV medical care.

**Conclusions/Importance:** Study findings highlight the need for increased scale-up of HIV testing efforts, as well as integrated HIV treatment and care in Kazakhstan.

### Keywords

HIV testing and treatment, injection drug use, Kazakhstan

### History

Received 10 September 2015

Revised 16 December 2015

Accepted 7 January 2016

### Introduction

Extensive research demonstrates that providing HIV testing and HIV medical care to key populations including people who inject drugs (PWID) comprise effective HIV prevention strategies (Dieffenbach & Fauci, 2009; Granich, Gilks, Dye, De Cock, & Williams, 2009; Mathers et al., 2010; Weinhardt, Carey, Johnson, & Bickham, 1999). Despite this evidence, coverage of HIV testing among PWID in Central Asia remains limited (Vickerman et al., 2014). Research indicates testing coverage in Kazakhstan to be 11.2% (Republican AIDS Center, 2015), while approximately 65% of PWID in Kazakhstan received an HIV test in the prior year and were aware of their status (Terlikbayeva et al., 2013). Among people who were HIV-positive, an estimated 65% were linked to care and 22–23% accessed antiretroviral therapy (ART), although this number may be lower among PWID who are HIV-positive (UNAIDS, 2012; Walsh & Maher, 2013).

Health care is provided free of charge for all citizens of Kazakhstan under the universally guaranteed package of services. HIV testing and syringe exchange services are included

in the guaranteed package and readily available at local polyclinics and non-governmental organizations (Republic of Kazakhstan, 2009). HIV diagnostics and treatment are also provided through government-run AIDS Centers free of charge, HIV testing in correctional settings is mandatory, and substance use treatment in narcological facilities is available free of charge for registered substance users. Free services available include detoxification, psychotherapy, rehabilitation, and therapeutic community living in government-run narcological centers (Ministry of Healthcare and Social Development, 2014). Opioid substitution therapy (OST) has been introduced as a pilot initiative in 10 sites across the country, but it is not available in Almaty city.

This paper examines 728 men and women who inject drugs and their intimate partners who participated in Project Renaissance, conducted between 2009 and 2012 in Almaty, Kazakhstan. We describe the prevalence of ever receiving an HIV test and ever-accessing HIV medical care. We examine how HIV testing and HIV medical care are associated with socio-demographic characteristics, access to general health-care and drug-related services (needle exchange services, drug treatment, and having a regular physician), and structural risk environment factors (homelessness, food insecurity, and ever being arrested and charged with a criminal offense).

Due to the role of social and structural factors in shaping contexts of HIV prevention and treatment access (Rhodes, Singer, Bourgois, Friedman, & Strathdee, 2005; Wolfe, Carrieri, & Shepard, 2010), we hypothesize that participants who have access to health- and drug-related services (needle exchange, drug treatment, and a regular physician) and who have ever been arrested are more likely to have received an HIV test and HIV medical care, while those who experience food insecurity and homelessness are less likely to have received HIV testing and HIV medical care.

## Method

### Study participants

We used baseline data from National Institute on Drug Abuse (NIDA)-funded Project Renaissance, conducted between 2009 and 2012 in Almaty, Kazakhstan. Study participants were recruited by trained research assistants through centers serving PWID, and word-of-mouth via other participants (El-Bassel et al., 2013). Participants were eligible to participate if they met the following criteria: (1) both were aged 18 or older; (2) both identified each other as their main partner of the opposite sex and someone whom the participant considered a boyfriend/girlfriend, spouse, lover and/or parent of his/her child; (3) both reported the relationship had existed for at least 3 months; (4) both reported intending to remain together for at least 12 months; (5) at least one partner reported having had unprotected vaginal or anal intercourse with the other partner within the previous 90 days; and (6) at least one partner reported injecting drugs within the previous 90 days. Participants were excluded if they met any of the following criteria: (1) either partner showed evidence of psychiatric, physical, or neurological impairment that would limit effective participation in the study; (2) either partner reported severe physical or sexual violence perpetrated by the other partner in the past year, as determined on a Revised Conflict Tactics Scale (Straus, 1979); (3) either partner reported that the couple was planning a pregnancy within the next 18 months; or (4) either partner was not fluent in Russian, as determined during the informed consent process.

Inclusion and exclusion criteria were based on needs for Project Renaissance, a randomized clinical trial of a HIV/hepatitis C/sexually transmitted infection prevention intervention, conducted with couples where one or both partners inject drugs (El-Bassel et al., 2014). A total of 971 individuals completed the screening interview. Of these, 728 met study eligibility criteria and completed the baseline interview. An additional 33 people were eligible but did not return to participate in the baseline. The Columbia University Institutional Review Board and the Kazakhstan School of Public Health Institutional Review Board approved all study procedures (El-Bassel et al., 2014).

### Data collection

Data collection included both self-reported data and objective biological assays. During the baseline visit, participants completed a 1.5-hour Audio Computer-Assisted Self-Interview (ACASI) conducted in Russian in a private room. The ACASI included all measures described below. After the

interview, a clinical research coordinator (CRC) conducted individual pre-test counseling, related to HIV, privately with each participant. Within 2 weeks of the baseline interview, the CRC notified each participant privately of his/her HIV test results, conducted post-test counseling, and provided referrals and navigation to treatment when applicable. Participants received the equivalent of USD\$10 for completing the baseline ACASI interview and biological testing.

## Measures

### Socio-demographic variables

Self-reported information was collected about participant's socio-demographic characteristics, including age, gender, years of education, ethnicity, marital status (legally married and common-law marriage as "married," and divorced, separated, widowed, or never married as "unmarried"), whether or not they had ever lived with their study partner, and whether participants had children.

### Service utilization

Participants were asked if they had ever utilized syringe exchange services, and whether they had ever attended drug treatment for heroin addiction. Additionally, participants were asked if they had a regular physician.

### HIV testing and HIV medical care

In the ACASI, participants reported whether they had ever received an HIV test. Participants who reported being HIV-positive in the ACASI were asked whether they ever received any medical care for HIV. The variable "medical care for HIV" thus refers to receiving any general medical care in relation to being HIV-positive. We also asked those who reported being HIV-positive in the ACASI whether they were taking ART.

### HIV status

HIV status was measured through the use of biological assays. A dried blood spot (DBS) technique was applied where a whole blood spot was obtained by a finger prick, applied to five printed circles on DBS filter paper cards, and sent to the reference laboratory at the Republican AIDS Center (RAC). For the serologic surveillance, a standard enzyme-linked immunosorbent assay (ELISA test manufactured by Abbott Murex) was used (Young, Moyes, Seagar, & McMillan, 1998). Tests were conducted using a serial two-test strategy, as recommended by the World Health Organization and routinely used at the RAC. US-manufactured Abbott Murex Biotech tests were used for the second test. According to the RAC Guidelines for Serological Surveillance, the Murex anti HIV ABBOTT has a reported sensitivity of >99.9% and specificity of 99%.

### Risk environment

We asked participants if they had been without a place to sleep in the past 90 days to measure homelessness, and

whether they had insufficient money for food in the past 90 days as an indicator of food insecurity. Additionally, we asked participants whether they had ever been arrested and charged with a criminal offense.

### Statistical analysis

Statistical analyses were performed in SPSS version 20. Bivariate relationships between receiving HIV testing; receiving HIV medical care; and each of the socio-demographic characteristics, service use, and risk environment factors were evaluated by *t*-tests for continuous variables and chi-square tests for categorical variables. Random effects logistic regression models were utilized to examine the relationship between HIV testing and HIV medical care outcomes with service utilization and risk environment factors, with couple identification number as a random effect to account for nesting of data within couples. HIV medical care outcomes were examined only among the sub-sample of participants who knew they were HIV-positive prior to study participation. Random effects models were conducted with a binomial probability distribution and logit link function. Each adjusted regression model included socio-demographic covariates (age, gender, ethnicity, marital status, and years of education). Unadjusted odds ratios (OR), adjusted odds ratios (aOR), and their associated 95% confidence intervals are reported.

## Results

### Socio-demographic characteristics

Participants' average age was 35 and most completed over 11 years of education (see Table 1). Nearly two-thirds of participants (65.7%) were ethnically Russian, although 11.7%

were Kazakh and 22.7% comprised another ethnic group (Ukrainian, German, Tatar, Uighur, Uzbek, Roma, Dungan, Korean, and others). Most respondents were married or in a common-law marriage (86.4%) and lived with their study partner (79.3%), while slightly more than half had children (52.9%). Of the sample, 96.4% of men ( $n = 351$ ) and 62.9% of women ( $n = 229$ ) reported ever injecting drugs (El-Bassel et al., 2013). Among the sub-sample of those who knew they were HIV-positive, 100% of men and 80.3% of women had injected drugs.

### Health and drug treatment service access and structural risk factors

Approximately one tenth of participants (11.1%) had ever utilized syringe exchange services, including 13.4% ( $n = 78$ ) of those who had ever injected drugs. A higher number (30.1%) reported ever attending drug treatment for heroin addiction, including 37.4% ( $n = 217$ ) of those who had ever injected drugs. A small number of participants who did not report ever injecting drugs reported utilizing syringe exchange services ( $n = 3$ ) and attending drug treatment for heroin addiction ( $n = 2$ ), suggesting perhaps that these few participants accessed drug treatment for non-injection heroin use, utilized needle exchange services for their partners or for other available services, or alternatively, that these participants did not accurately report their own history of injection drug use. Over one-fourth (27.5%) of the sample had a regular physician.

Examining the structural risk environment, 13.5% of participants had experienced homelessness in the prior 90 days, and nearly half of participants (48.8%) experienced food insecurity during the same period. Two-thirds (67.0%) of

Table 1. Socio-demographic characteristics, service use, and exposure to risk environment by HIV testing ( $N = 728$ ).

	Total sample ( $N = 728$ )	Ever had an HIV test ( $n = 544$ )	Never had an HIV test ( $n = 184$ )
<b>Socio-demographics</b>	35.75, 7.8	36.0, 7.6	35.0, 8.4
Age (mean, SD)			
Years of education completed (mean, SD)	11.4, 3.3	11.4, 3.2	11.5, 3.5
<b>Gender (<math>n, \%</math>)</b>			
Male	364 (50)	275 (50.6)	89 (48.4)
Female	364 (50)	269 (49.4)	95 (51.6)
<b>Ethnicity</b>			
Kazakh	85 (11.7)	63 (11.6)	22 (11.9)
Russian	478 (65.7)	357 (65.6)	121 (65.8)
Other	165 (22.7)	124 (22.8)	41 (22.3)
<b>Marital status</b>			
Married	629 (86.4)	471 (86.6)	158 (85.9)
Unmarried	99 (13.6)	73 (13.4)	26 (14.1)
Ever lived with study partner	577 (79.3)	432 (79.4)	145 (78.8)
Has children	385 (52.9)	301 (55.3)*	84 (45.7)*
<b>Access to services</b>			
Ever utilized syringe exchange services	81 (11.1)	70 (12.9)*	11 (6.0)*
Ever attended drug treatment for heroin addiction	219 (30.1)	189 (34.7)**	30 (16.3)**
Has a regular physician	200 (27.5)	162 (29.8)*	38 (20.7)*
<b>Structural risk environment</b>			
Homeless in past 90 days	98 (13.5)	66 (12.1)	32 (17.4)
Food insecurity in past 90 days	355 (48.8)	270 (49.6)	85 (46.2)
Ever arrested and charged with a criminal offense	488 (67.0)	393 (72.2)**	95 (51.6)**

Note: The significance test for differences was performed by *t*-test and chi-square test.

\* $p < 0.05$ , \*\* $p < 0.01$

study participants had ever been arrested and charged with a criminal offense.

### HIV testing and socio-demographic characteristics, access to services, and structural risk environment

Of 728 study participants, 544 (74.7%) had received an HIV test. Table 1 examines bivariate relationships between HIV testing and socio-demographic characteristics, service usage, and structural risk factors. No significant differences in age, years of education, ethnicity, marital status, or ever living with one's study partner were found in respect to HIV testing. Participants with children were more likely to have been tested for HIV than those without children ( $p < .05$ ). Those who utilized syringe exchange services, attended drug treatment for heroin addiction, or had a regular physician were more likely to have been tested when compared to those who had lower levels of service access (see Table 1). Additionally, those who were arrested and charged with a criminal offense were more likely to have been tested than those who were not been arrested. Food insecurity and homelessness were not associated with HIV testing.

### HIV medical treatment and socio-demographic characteristics, access to services, and structural risk environment

Although 183 participants (25.1%) tested HIV-positive, 141 (77.0%) of those who tested positive were previously aware of their HIV-positive status. Of these individuals, 73 (49.6%) ever accessed HIV medical care. Few ( $n = 22$ ) were taking ART, representing 15.6% of those who were aware of their HIV-positive status (El-Bassel et al., 2013).

When analyzing access to HIV medical care, this paper focuses only on the sub-sample who knew they were HIV-positive ( $n = 141$ ). Similar to findings regarding HIV testing, socio-demographic characteristics did not vary significantly by whether participants accessed HIV medical care, although participants with children were again more likely to have accessed HIV medical care ( $p < .05$ ; see Table 2). Participants who utilized syringe exchange services were more likely to have received HIV medical care ( $p < .05$ ), as were those had a regular physician ( $p < .001$ ). People who experienced food insecurity were less likely to have accessed medical care than those without food insecurity ( $p < .05$ ).

### Multivariate analysis of HIV testing and HIV medical care

We examined study hypotheses that participants who utilized needle exchange services, attended heroin treatment, accessed a regular doctor, or were ever arrested were (1) more likely to be tested for HIV and (2) more likely to receive HIV medical care; and the hypothesis that participants who experienced food insecurity or homelessness were (1) less likely to receive HIV testing and (2) less likely to receive HIV medical care, when controlling for socio-demographic characteristics.

Overall, unadjusted and adjusted models produced similar measures of association; adjusted models are reported here. Accessing HIV testing was associated with utilizing needle

exchange services (aOR = 2.219, 95% CI = 1.122, 4.384), attending drug treatment for heroin addiction (aOR = 2.804, 95% CI = 1.793, 4.384), and having a regular physician (aOR = 1.685, 95% CI = 1.105, 2.570) in support of our hypotheses (see Table 3). HIV testing was not associated with homelessness and food insecurity. The odds of having an HIV test among those who were ever arrested and charged with a criminal offense was 2.721 times the odds of those who were not arrested in the adjusted model (95% CI = 1.820–4.067).

Receiving HIV medical care was associated with utilizing needle exchange services (aOR = 3.456, 95% CI = 1.092, 10.892) and having a regular physician (aOR = 4.884, 95% CI = 2.032, 11.752), but not with attending drug treatment (aOR = 1.177, 95% CI = .462, 3.001) (see Table 4). In the multivariate models, receiving HIV medical care was not associated with risk environment characteristics, including criminal justice involvement.

## Discussion

Study findings suggest that access to HIV testing and HIV medical care among PWID and their intimate partners in Kazakhstan remains limited, where three quarters of our sample ever had an HIV test, and only half of those who knew they were HIV-positive ever accessed general HIV-related medical care. Coverage of testing reflects scale-up efforts and may be higher than that found in other Central Asian countries (Thorne, Ferencic, Malyuta, Mimica, & Niemiec, 2010). Similarly, while access to HIV medical care was low, our finding that 15.6% of HIV-positive participants accessed ART demonstrates higher levels of access than that observed in Uzbekistan, Russia, Pakistan, Chile, and Kenya, where less than 1% of PWID received ART (Mathers et al., 2010).

Study hypotheses that testing and treatment were higher among participants accessing needle exchange, drug treatment, and general health-care services were partially supported. Accessing needle exchange services and having a regular physician were associated with receiving HIV testing and receiving HIV medical care, perhaps due to the frequent co-location of needle exchange services within HIV general polyclinics or AIDS Centers, where general medical care is also available. Syringe exchange programs may be an important venue for the promotion of HIV prevention and access to treatment (Vlahov, 2010), although additional research is needed to determine pathways of service access.

Drug treatment for heroin was associated with HIV testing, but not with HIV medical care, suggesting drug treatment services may be a missed opportunity for linking HIV-positive individuals to necessary medical care and treatment. Research suggests that opiate substitution therapies, particularly methadone but also buprenorphine and naltrexone, lead to improvements in adherence to ART and subsequent reductions in the likelihood of HIV transmission (Malta, Magnanini, Strathdee, & Bastos, 2010; Metzger, Woody, & O'Brien, 2010). Provision of evidence-based drug treatment services alongside clinical treatment for HIV is feasible (Altice et al., 2011), and this could be a useful approach to

Table 2. Socio-demographic characteristics, service use, and exposure to risk environment by access to HIV medical care among people aware of HIV-positive status ( $N = 141$ ).

	Total sample ( $N = 141$ )	Ever had access to HIV medical care ( $n = 70$ )	No access ( $n = 71$ )
<b>Socio-demographics</b>			
Age	36.5 (7.4)	36.9 (8.1)	36.1 (6.6)
Years of education completed (mean, SD)	11.1 (3.7)	10.7 (2.4)	11.4 (4.7)
<b>Gender (n, %)</b>			
Male	80 (56.7)	38 (54.3)	42 (59.2)
Female	61 (43.3)	32 (45.7)	29 (40.8)
<b>Ethnicity</b>			
Kazakh	12 (8.5)	3 (4.3)	9 (12.7)
Russian	98 (69.5)	47 (67.1)	51 (71.8)
Other	31 (22.0)	20 (28.6)	11 (15.5)
<b>Marital Status</b>			
Married	119 (84.4)	62 (88.6)	57 (80.3)
Unmarried	22 (15.6)	8 (11.4)	14 (19.7)
Ever lived with study partner	116 (82.3)	59 (84.3)	57 (80.3)
Has children	73 (51.8)	40 (57.1)*	33 (46.5)*
<b>Access to services</b>			
Ever utilized syringe exchange services	27 (19.1)	19 (27.1)*	8 (11.3)*
Ever attended drug treatment for heroin addiction	48 (34.0)	24 (34.3)	24 (33.8)
Has a regular physician	68 (48.2)	46 (65.7)**	22 (31.0)**
<b>Structural risk environment</b>			
Homeless in past 90 days	22 (15.6)	11 (15.7)	11 (15.5)
Food insecurity in past 90 days	67 (47.5)	26 (37.1)*	41 (57.7)*
Ever arrested and charged with a criminal offense	121 (85.8)	58 (82.9)	63 (88.7)

\* $p < 0.05$ , \*\* $p < .01$ .Table 3. Mixed-effect logistic regression of predictors of HIV testing with random effects for couple ( $N = 728$ )<sup>a</sup>.

	Unadjusted	Adjusted
<b>Access to services</b>		
Ever utilized needle exchange services	2.307 (1.176, 4.527)*	2.219 (1.122, 4.384)*
Ever attended drug treatment for heroin addiction	2.694 (1.742, 4.170)**	2.804 (1.793, 4.384)**
Has a regular physician	1.616 (1.070, 2.440)*	1.685 (1.105, 2.570)*
<b>Structural risk environment</b>		
Homeless in the past 90 days	0.644 (0.399, 1.040)	0.643 (0.395, 1.048)
Insufficient money for food in past 90 days	1.150 (0.813, 1.627)	1.133 (0.798, 1.611)
Ever arrested and charged with a criminal offense	2.406 (1.692, 3.421)**	2.721 (1.820, 4.067)**

<sup>a</sup>Each model controlled for age, gender, ethnicity, marital status, and years of education.\* $p < 0.05$ , \*\* $p < 0.01$ Table 4. Mixed-effect logistic regression of predictors of receiving HIV medical care with random effects for couple among people aware of HIV-positive status ( $n = 141$ )<sup>a</sup>.

	Unadjusted	Adjusted
<b>Access to services</b>		
Ever utilized needle exchange services	3.025 (1.047, 8.732)*	3.456 (1.092, 10.892)*
Ever attended drug treatment for heroin addiction	0.959 (0.424, 2.166)	1.177 (0.462, 3.001)
Has a regular physician	4.071 (1.833, 9.043)**	4.884 (2.032, 11.752)**
<b>Structural risk environment</b>		
Homeless in the past 90 days	1.094 (0.374, 3.203)	1.066 (0.346, 3.281)
Insufficient money for food in past 90 days	.463 (0.213, 1.008)	.493 (0.215, 1.134)
Ever arrested and charged with criminal offense	.588 (0.193, 1.791)	.533 (0.145, 1.964)

<sup>a</sup>Each model controlled for age, gender, ethnicity, marital status, and years of education.\* $p < 0.05$ , \*\* $p < 0.01$ 

improving both drug treatment and HIV-related services for people who use drugs in Kazakhstan.

The association observed between HIV testing and criminal justice involvement likely coincides with mandatory

testing policies within the prison system (Thorne et al., 2010). However, we did not find any association between accessing HIV medical care and being arrested and charged with a criminal offense. While the criminal justice system is

an avenue within which testing and treatment can be delivered, access to care within prison systems is often negligible, and links to care upon release are necessary (Azbel, Wickersham, Grishaev, Dvoryak, & Altice, 2013).

Contrary to study hypotheses, homelessness and food insecurity were not associated with decreased likelihood of accessing HIV testing or HIV medical care in the multivariate models, suggesting that services reached those from various socio-economic backgrounds. The bivariate finding that participants with children were more likely to be tested and access HIV medical care may suggest that parents are more connected to health systems and medical care, perhaps due to mandatory HIV testing and subsequent treatment when needed during pregnancy.

Although scaling up HIV prevention services including testing, drug treatment, and ART can avert deaths and significantly reduce health-care costs (Schwartländer et al., 2011), access to these services has been found to be low worldwide, especially among PWID (Degenhardt et al., 2010; Niccolai et al., 2010). People who are HIV-positive and unaware of their status tend to change their risk behaviors upon being informed of their status (Marks, Crepaz, Senterfitt, & Janssen, 2005), highlighting the need for access to regular HIV testing among those with high-risk behavior. To reduce HIV incidence, coverage of cost-effective measures including HIV testing and counseling, needle exchange programs, medication-assisted therapy, and ART needs to reach at least 60% (Dutta, Wirtz, Baral, Beyrer, & Cleghorn, 2012). UNAIDS's (Joint United Nations Programme on HIV and AIDS) ambitious treatment target is that, by 2020, 90% of all people living with HIV will know their HIV status, 90% of people with diagnosed HIV infection will receive sustained ART, and 90% of those receiving ART will have viral suppression (UNAIDS, 2014).

Working toward these treatment targets in Kazakhstan, additional efforts to engage communities of people who inject drugs are needed. As HIV testing is provided only within government-run facilities in Kazakhstan, further scale-up of HIV testing may be possible through community-driven HIV testing and counseling efforts, which have demonstrated promise globally (Suthar et al., 2013). The City AIDS Center in Almaty, where study participants with positive HIV tests were referred, provided HIV clinical services, including laboratory monitoring and case management of HIV-positive patients by a multidisciplinary team of specialists (Ministry of Health, 2011). However, linkages across service sectors such as drug treatment and co-occurring infections, i.e., TB and HCV, were limited. Removing structural barriers to the integration of harm reduction and HIV treatment services is critical for the promotion of combination intervention approaches across Central Asia (Jolley et al., 2012).

Availability of preventative HIV health services may be insufficient to ensure access to testing and treatment when additional barriers including stigma, criminalization, discriminatory practices, and limited knowledge persist (Spicer et al., 2011). These barriers may lead to reluctance among PWID to receive HIV testing from government-run services, even when services are anonymous (Thorne, 2010). Research globally has identified the importance of coverage of a combination of services, including syringe exchange, OST, and

ART, alongside attention to social and structural barriers (Degenhardt et al., 2010) in order to prevent HIV infection. Continued attention to social and structural barriers to the provision of accessible combined approaches in Kazakhstan is needed (Krüsi, Wood, Montaner, & Kerr, 2010).

A primary limitation of the study is that the sample is not random, thus findings do not apply to all PWID in Kazakhstan. Furthermore, data was cross-sectional, and we cannot determine how access to services and structural risk environment factors may have influenced access to HIV testing and treatment. For example, while we examined having a regular physician and accessing drug treatment as independent correlates of HIV testing and HIV medical care, persons who test positive for HIV may be more likely to seek out such services. We were also limited in our ability to examine HIV treatment outcomes. We measured HIV medical care as a general variable rather than specifically looking at ART. However, a regular physician may have provided HIV medical care, and participants may not have distinguished clearly between HIV medical care and medical care received for other conditions. Although our sample includes a portion of participants who did not report injecting drugs, we chose to include these non-injecting intimate partners in the analysis as they also experience risks for HIV and are in need of access to HIV testing and HIV medical care (Jenness, Neaigus, Hagan, Murrill, & Wendel, 2010). Partner and familial supports may be a relevant social component of treatment access for many PWID (Krüsi et al., 2010). We encourage further research into the role of relationship characteristics and social support in treatment promotion, as well as to the service needs of intimate partners of PWID (Hammett, Van, Kling, Binh, & Oanh, 2010; Solomon, Mehta, Latimore, Srikrishnan, & Celentano, 2010).

## Conclusions

Study findings suggest that access to HIV testing is higher among PWID and their intimate partners who access syringe exchange services, drug treatment, a regular physician, and the criminal justice system. While testing has reached a large proportion of PWID and their intimate partners, additional attention is needed to increase access to regular testing and counseling for couples where one or both partners inject in Kazakhstan. Among people who knew they were HIV-positive, those who received syringe exchange services and had a regular physician had higher rates of access to general HIV medical treatment. While links or co-location between services appear to be helpful, increased integration across HIV, drug treatment, and criminal justice systems are needed.

## Acknowledgments

We would like to thank each participant who shared her or his time and experience with this project, and would also like to acknowledge the project staff members who assisted in recruiting participants and conducting assessments.

## Funding

This study was supported by the National Institute of Drug Abuse # R01 DA022914-01A2 to Dr. Nabila El-Bassel.

## Declaration of interest

The authors have no conflict of interest to report.

## References

- Alice, F. L., Bruce, R. D., Lucas, G. M., Lum, P. J., Korhuis, P. T., Flanagan, T. P., ... Fiellin, D. A. (2011). HIV treatment outcomes among HIV-infected, opioid-dependent patients receiving buprenorphine/naloxone treatment within HIV clinical care settings: Results from a multisite study. *JAIDS*, 56(Suppl 1), S22–S32.
- Azbel, L., Wickersham, J. A., Grishaev, Y., Dvoryak, S., & Altice, F. L. (2013). Burden of infectious diseases, substance use disorders, and mental illness among Ukrainian prisoners transitioning to the community. *PLoS ONE*, 8(3), 1–9.
- Degenhardt, L., Mathers, B., Vickerman, P., Rhodes, T., Latkin, C., & Hickman, M. (2010). Prevention of HIV infection for people who inject drugs: Why individual, structural, and combination approaches are needed. *The Lancet*, 376(9737), 285–301.
- Dieffenbach, C. W., & Fauci, A. S. (2009). Universal voluntary testing and treatment for prevention of HIV transmission. *JAMA*, 301(22), 2380–2382.
- Dutta, A., Wirtz, A. L., Baral, S., Beyrer, C., & Cleghorn, F. R. (2012). Key harm reduction interventions and their impact on the reduction of risky behavior and HIV incidence among people who inject drugs in low-income and middle-income countries. *Current Opinion in HIV/AIDS*, 7, 362–368.
- El-Bassel, N., Gilbert, L., Terlikbayeva, A., Wu, E., Beyrer, C., Shaw, S., ... Tukeyev, M. (2013). HIV among injection drug users and their intimate partners in Almaty, Kazakhstan. *AIDS and Behavior*, 17(7), 2490–2500.
- El-Bassel, N., Gilbert, L., Terlikbayeva, A., Beyrer, C., Wu, E., Chang, M., ... Rozental, Y. (2014). Effects of a couple-based intervention to reduce risks for HIV, HCV, and STIs among drug-involved heterosexual couples in Kazakhstan: a randomized controlled trial. *Journal of Acquired Immune Deficiency Syndromes*, 67(2), 196–203.
- Granich, R. M., Gilks, C. F., Dye, C., De Cock, K. M., & Williams, B. G. (2009). Universal voluntary HIV testing with immediate antiretroviral therapy as a strategy for elimination of HIV transmission: A mathematical model. *The Lancet*, 373(9657), 48–57.
- Hammett, T. M., Van, N. T. H., Kling, R., Binh, K. T., & Oanh, K. T. Hai. (2010). Female sexual partners of injection drug users in Vietnam: An at-risk population in urgent need of HIV prevention services. *AIDS Care*, 22(12), 1466–1472.
- Jenness, S. M., Neaigus, A., Hagan, H., Murrill, C. S., & Wendel, T. (2010). Heterosexual HIV and sexual partnerships between injection drug users and noninjection drug users. *AIDS Patient Care and STDs*, 24(3), 175–181.
- Krüsi, A., Wood, E., Montaner, J., & Kerr, T. (2010). Social and structural determinants of HAART access and adherence among injection drug users. *International Journal of Drug Policy*, 21(1), 4–9.
- Malta, M., Magnanini, M. M. F., Strathdee, S. A., & Bastos, F.I. (2010). Adherence to antiretroviral therapy among HIV-infected drug users: A meta-analysis. *AIDS and Behavior*, 14(4), 731–747.
- Marks, G., Crepaz, N., Senterfitt, J. W., & Janssen, R. S. (2005). Meta-analysis of high-risk sexual behavior in persons aware and unaware they are infected with HIV in the United States: Implications for HIV prevention programs. *Journal of Acquired Immune Deficiency Syndromes (1999)*, 39(4), 446–453.
- Mathers, B. M., Degenhardt, L., Ali, H., Wiessing, L., Hickman, M., Mattick, R. P., ... Injecting, Drug. (2010). HIV prevention, treatment, and care services for people who inject drugs: A systematic review of global, regional, and national coverage. *The Lancet*, 375, 1014–1028.
- Metzger, David S, Woody, George E, & O'Brien, Charles P. (2010). Drug treatment as HIV prevention: A research update. *JAIDS*, 55 (Suppl 1), S32–S32.
- Niccolai, L. M., Tousseva, O. V., Verevokhin, S. V., Barbour, R., Heimer, R., & Kozlov, A. P. (2010). High HIV prevalence, suboptimal HIV testing, and low knowledge of HIV-positive serostatus among injection drug users in St. Petersburg, Russia. *AIDS and Behavior*, 14 (4), 932–941.
- Rhodes, T., Singer, M., Bourgois, P., Friedman, S. R., & Strathdee, S.A. (2005). The social structural production of HIV risk among injecting drug users. *Social Science & Medicine*, 61(5), 1026–1044.
- Schwartzländer, B., Stover, J., Hallett, T., Atun, R., Avila, C., Gouws, E., ... Barr, D. (2011). Towards an improved investment approach for an effective response to HIV/AIDS. *The Lancet*, 377 (9782), 2031–2041.
- Solomon, S. S., Mehta, S. H., Latimore, A., Srikrishnan, A. K., & Celentano, D. D. (2010). The impact of HIV and high-risk behaviours on the wives of married men who have sex with men and injection drug users: Implications for HIV prevention. *Journal of the International AIDS Society*, 13(Suppl 2), 1–8.
- Spicer, N., Bogdan, D., Brugha, R., Harmer, A., Murzalieva, G., & Semigina, T. (2011). "It's risky to walk in the city with syringes": Understanding access to HIV/AIDS services for injecting drug users in the former Soviet Union countries of Ukraine and Kyrgyzstan. *Growth*, 1, 1–15.
- Straus, M. A. (1979). Measuring intrafamily conflict and violence: The conflict tactics (CT) scales. *Journal of Marriage and the Family*, 41(1), 75–88.
- Suthar, A. B., Ford, N., Bachanas, P. J., Wong, V. J., Rajan, J. S., Saltzman, A. K., ... Negussie, E. K. (2013). Towards universal voluntary HIV testing and counselling: A systematic review and meta-analysis of community-based approaches. *PLoS Medicine*, 10(8), 1–23.
- Terlikbayeva, A., Zhussupov, B., Primbetova, S., Gilbert, L. Atabekov, N., Giyasova, G., ... El-Bassel, N. (2013). Access to HIV counseling and testing among people who inject drugs in Central Asia: Strategies for improving access and linkages to treatment and care. *Drug and Alcohol Dependence*, 132, S61–S64.
- Thorne, C., Ferencic, N., Malyuta, R., Mimica, J., & Niemiec, T. (2010). Central Asia: Hotspot in the worldwide HIV epidemic. *The Lancet Infectious Diseases*, 10(7), 479–488.
- UNAIDS. (2012). Global Report: UNAIDS Report on the Global AIDS Epidemic 2012. Switzerland: Joint United Nations Programme on HIV/AIDS (UNAIDS).
- Vickerman, P., Platt, L., Jolley, E., Rhodes, T., Kazatchkine, M.D., & Latypov, A. (2014). Controlling HIV among people who inject drugs in Eastern Europe and Central Asia: Insights from modeling. *International Journal of Drug Policy*, 25(6), 1163–1173.
- Walsh, N., & Maher, L. (2013). HIV and HCV among people who inject drugs in Central Asia. *Drug and Alcohol Dependence*, 132, S37–S40.
- Weinhardt, L. S., Carey, M. P., Johnson, B. T., & Bickham, N. L. (1999). Effects of HIV counseling and testing on sexual risk behavior: A meta-analytic review of published research, 1985–1997. *American Journal of Public Health*, 89(9), 1397–1405.
- Wolfe, D., Carrieri, M. P., & Shepard, D. (2010). Treatment and care for injecting drug users with HIV infection: A review of barriers and ways forward. *The Lancet*, 376(9738), 355–366.
- Young, H., Moyes, A., Seagar, L., & McMillan, A. (1998). Novel recombinant-antigen enzyme immunoassay for serological diagnosis of syphilis. *Journal of Clinical Microbiology*, 36(4), 913–917.